

Chern numbers and localization by non-Hermitian operators

Jonathan Miller¹ and Peter B. Weichman²

¹*NEC Research Institute, 4 Independence Way, Princeton, NJ 08540*

²*Blackhawk Geometrics, 301 Commercial Rd., Suite B, Golden, CO 80401*

(February 1, 2008)

In Hermitian quantum mechanics, extended current-carrying states are distinguished from localized ones by their non-zero Chern number. We generalize the notion of Chern number to non-Hermitian localization problems such as tilted flux lines in type-II superconductors with line defects and passive scalar transport in random vorticity fields. We find that the usual point eigenvalue degeneracies that occur in the Hermitian case are now loops bounding a branch cut sheet with the same quantized total Berry flux, and hence Chern number, as the original point. If a loop cuts the integration surface non-quantized contributions to the total flux result.

In Hermitian quantum mechanics the differential geometric notion of Chern number arises in the context of adiabatic transport of eigenstates in Hilbert space [1]. The change in the overall phase of a state after it is transported around a closed loop contains a purely geometric contribution associated with curvature of the Hilbert space. Quantized integer Chern numbers are associated with singularities in the curvature that originate in eigenvalue degeneracies of the Hamiltonian generating the transport.

One way of transporting an eigenstate is by changing its boundary conditions. A Chern number can then be computed as a certain integral over the space of boundary conditions. Since a localized state is insensitive to boundary conditions, a nonzero Chern number should be associated only with an extended state, and serves as a nontrivial diagnostic of localization. This procedure has been used to study localization in the integer quantum Hall effect [2]. Physically, the Chern number reflects the ability of the state to transport charge across the system, and is proportional to its Hall conductance [1].

Recently there has been much interest in localization properties of certain *non*-Hermitian operators, for example, operators governing effective two-dimensional quantum descriptions of pinning of tilted flux lines by extended defects in three-dimensional type-II superconductors [3], and passive scalar transport by two-dimensional random vorticity fields [4]. Both these examples have close ties to the problem of electron localization in random magnetic fields [6]. In this paper we will consider the generalization of Chern numbers to non-Hermitian operators. We review the non-Hermitian Berry phase [5] and how it gives rise to Chern numbers through eigenvalue degeneracies. Application is then made to a class of operators that interpolate, as a function of a parameter $0 \leq \theta < \pi/2$, between a special case of the Hermitian random flux model and the non-Hermitian flux line and passive scalar problems.

Consider a family $\mathbf{L}(\mathbf{X})$ of (possibly non-Hermitian) operators, where $\mathbf{X} = (X_1, X_2, \dots)$ is a set of fixed *real*

parameters spanning a manifold. Let $|m; \mathbf{X}\rangle$ and $\langle m; \mathbf{X}|$ be the right and left eigenvectors of $\mathbf{L}(\mathbf{X})$ satisfying

$$\begin{aligned} \mathbf{L}(\mathbf{X})|m; \mathbf{X}\rangle &= E_m(\mathbf{X})|m; \mathbf{X}\rangle \\ \langle m; \mathbf{X}|\mathbf{L}(\mathbf{X}) &= E_m(\mathbf{X})\langle m; \mathbf{X}| \\ \langle m; \mathbf{X}|n; \mathbf{X}\rangle &= \delta_{mn}. \end{aligned} \quad (1)$$

The eigenstates are determined up to an arbitrary complex amplitude, and we make a fixed arbitrary choice of this quantity so that $|m; \mathbf{X}\rangle$ is a single-valued function of $\mathbf{X}$ over the manifold. The projection operator, $\mathbf{P}_m(\mathbf{X}) = |m; \mathbf{X}\rangle\langle m; \mathbf{X}|$, is independent of this amplitude. The eigenvalues $E_m(\mathbf{X})$, assumed nondegenerate for now, may be complex, and in general $\langle m; \mathbf{X}|^\dagger \neq |m; \mathbf{X}\rangle$. For any given operator $\mathbf{L}(t)$, we define the dynamics of states $|\phi(t)\rangle$ and $\langle\phi(t)|$ via

$$i\partial_t|\phi(t)\rangle = \mathbf{L}(t)|\phi(t)\rangle, \quad -i\partial_t\langle\phi(t)| = \langle\phi(t)|\mathbf{L}(t). \quad (2)$$

This dynamics preserves all inner products $\langle\phi_1(t)|\phi_2(t)\rangle$.

Let $\mathbf{X}(\alpha)$, $0 \leq \alpha \leq 1$, $\mathbf{X}(0) = \mathbf{X}(1)$, parameterize a closed path, C , in parameter space. We define *parallel transported* eigenstates $|m; \alpha\rangle = b_m(\alpha)|m; \mathbf{X}(\alpha)\rangle$ along the path $\mathbf{X}(\alpha)$ by the rule

$$\partial_\alpha b_m(\alpha) = -\langle m; \mathbf{X}(\alpha)|\partial_\alpha |m; \mathbf{X}(\alpha)\rangle b_m(\alpha), \quad (3)$$

with $b_m(0) = 1$. The solution is [5]

$$\begin{aligned} b_m(\alpha) &= \exp \left[i \int_{C: \mathbf{X}(0)}^{\mathbf{X}(\alpha)} \mathbf{A}_m(\mathbf{X}) \cdot d\mathbf{X} \right] \\ \mathbf{A}_m(\mathbf{X}) &\equiv \langle m; \mathbf{X}(\alpha)|i\partial_\mathbf{X}|m; \mathbf{X}(\alpha)\rangle. \end{aligned} \quad (4)$$

Direct substitution shows that this parallel transport may be generated dynamically using the operator $\mathbf{L}(t) = i[\partial_t \mathbf{P}_m(\mathbf{X}(t)), \mathbf{P}_m(\mathbf{X}(t))]$ (with $\alpha = t$) [7]. For a closed loop one obtains

$$b_m(1) = e^{i\gamma_m(C)}, \quad \gamma_m(C) = \int_C d\mathbf{X} \cdot \mathbf{A}_m(\mathbf{X}). \quad (5)$$

In the Hermitean case $\mathbf{A}_m$, and hence $\gamma(C)$, is real. In the non-Hermitean case they are both generally complex. It is easy to see that $\gamma(C)$ is independent of the arbitrary normalization of the states $|m; \mathbf{X}\rangle$. Any change of normalization, $e^{i\lambda(\mathbf{X})}$, appears as a gauge transformation $\mathbf{A}' = \mathbf{A} - \partial_{\mathbf{X}}\lambda$, under which the integral in (5) is invariant. Provided no singularities are encountered we may write $\gamma(C)$ as the integral of the gauge invariant flux using Stokes theorem,

$$\begin{aligned} \mathbf{F}_m(\mathbf{X}) &= \nabla \times \mathbf{A}_m(\mathbf{X}) \\ &= i[\partial_{\mathbf{X}}\langle m_0; \mathbf{X} | \times [\partial_{\mathbf{X}} | m_0; \mathbf{X} \rangle], \end{aligned} \quad (6)$$

over any surface S bounded by C [8]:

$$\gamma_m(C) = \int_S d\Sigma \cdot \mathbf{F}_m(\mathbf{X}). \quad (7)$$

By construction one has $\nabla \cdot \mathbf{F}_m = 0$, as is required for $\gamma(C)$ to be independent of the choice of S .

By inserting a complete set of states, the flux may be written in the form

$$\mathbf{F}_{m_0}(\mathbf{X}) = i \sum_n \frac{\langle m; \mathbf{X} | \partial_{\mathbf{X}} \mathbf{L} | n; \mathbf{X} \rangle \times \langle n; \mathbf{X} | \partial_{\mathbf{X}} \mathbf{L} | m; \mathbf{X} \rangle}{(E_n - E_m)^2} \quad (8)$$

where the prime indicates that the term $n = m$ is omitted. Equation (8) demonstrates explicitly that so long as $\mathbf{L}(\mathbf{X})$ is smooth, singularities in $\mathbf{F}_m$ can only occur when two eigenvalues cross: $E_n - E_m \rightarrow 0$ for some $n \neq m$. In the Hermitean case eigenvalue crossings, and therefore singularities in $\mathbf{F}_m(\mathbf{X})$, generically occur at isolated points in the three dimensional space of $\mathbf{X}$. One may choose the surface S in (7) to avoid such points by passing either over or under them. On the one hand, the difference of the integrals over two such surfaces S_1 and S_2 is

$$\gamma_m(S) \equiv \gamma_m(S_1) - \gamma_m(S_2) = \int_S d\Sigma \cdot \mathbf{F}_m, \quad (9)$$

where $S = S_1 \cup S_2$ is a closed surface enclosing the singularity. On the other hand, $e^{i\gamma_m(S_1)} = e^{i\gamma_m(S_2)}$ since both surfaces are bounded by the same curve C . We conclude that for any closed surface S , $\gamma_m(S) = 2\pi q$ for some integer *Chern number* q [9]. Eigenvalue degeneracies act as a quantized point sources of flux for $\mathbf{F}_m$.

This argument does not rely directly on Hermiticity, but only on the result that energy level degeneracies occur generically at points in three dimensional space, so that these singularities can be enclosed by singularity-free closed surfaces. We now address the issue of how to generalize this result to the non-Hermitean case. In order to study a degeneracy in more detail, we project $\mathbf{L}$ onto the (complex) two-dimensional subspace spanned by the eigenvectors $|m; \mathbf{X}\rangle$ and $|n; \mathbf{X}\rangle$. If $\mathbf{P}_{mn} = |m\rangle\langle m| + |n\rangle\langle n|$ is the corresponding orthogonal projection, we consider the matrix

$$\begin{aligned} \mathbf{L}_{mn}(\mathbf{X}) &= \mathbf{P}_{mn}(\mathbf{X})\mathbf{L}(\mathbf{X})\mathbf{P}_{mn}(\mathbf{X}) \\ &= \alpha_0(\mathbf{X})\mathbf{I} + \vec{\alpha}(\mathbf{X}) \cdot \vec{\sigma} \end{aligned} \quad (10)$$

where $\vec{\sigma}$ are the Pauli matrices, and α_0 and $\vec{\alpha} = (\alpha_1, \alpha_2, \alpha_3)$ are four (at this stage arbitrary) complex numbers. In the Hermitean case these numbers are all real. The eigenvalues of $\mathbf{L}_{mn}$ are $E_{\pm} = \alpha_0 \pm \Delta E$, where $\Delta E = \sqrt{\vec{\alpha} \cdot \vec{\alpha}}$. The left and right eigenvectors are easily computed as well, and the fluxes are

$$\mathbf{F}_{\pm}(\mathbf{X}) = \mp \frac{\alpha_1 \mathbf{a}_2 \times \mathbf{a}_3 + \alpha_2 \mathbf{a}_3 \times \mathbf{a}_1 + \alpha_3 \mathbf{a}_1 \times \mathbf{a}_2}{2\Delta E^3}, \quad (11)$$

where $\mathbf{a}_i(\mathbf{X}) = \partial_{\mathbf{X}}\alpha_i(\mathbf{X})$.

Equation (11) can be used to explore the nature of the singularities in $\mathbf{F}_{\pm}$ in the neighborhood of degeneracies, $\Delta E \rightarrow 0$. The condition for $\Delta E = 0$ is simply $\alpha_1^2 + \alpha_2^2 + \alpha_3^2 = 0$. In the Hermitean case where the α_i are all real, this condition requires that $\alpha_1 = \alpha_2 = \alpha_3 = 0$, but in general this equality places only *two* conditions on the six parameters (the real and imaginary parts of each α_i , $i = 1, 2, 3$). In the parameter space $\mathbf{X}$ degeneracies will then generically occur on a submanifold of codimension 2, i.e., on *lines* in the three dimensional space.

Since the matrix $\mathbf{L}_{mn}$ is no longer Hermitean, there is no guarantee that there will be two independent eigenvectors associated with the two degenerate eigenvalues. Generically $\mathbf{L}_{mn}$ will be upper-triangular (with $\alpha_1 + i\alpha_2 = 0$ at the same time that $\alpha_3 = 0$, but $\alpha_1 - i\alpha_2 \neq 0$), and only one eigenvector will exist, with the orthogonal direction spanned by a *generalized eigenvector* [10]. We shall call this situation a *generalized degeneracy*. What we call a *true degeneracy* exists only when both $\alpha_1 \pm i\alpha_2 = 0$, so that $\mathbf{L}_{mn} = E\mathbf{I}$ is proportional to the identity matrix (this differs from the Hermitean case only in that $\alpha_0 = E$ may be complex), and places *six* conditions on the occurrence of a true degeneracy, which will generically *not* occur for a three dimensional parameter space $\mathbf{X}$, unless further restrictions are placed on the class of matrices (such as self-adjointness). In what follows we do not impose any such further restrictions.

If $\mathbf{X}_g$ be a point of generalized degeneracy, set $\mathbf{x} = \mathbf{X} - \mathbf{X}_g$. From (11) we obtain for small $\mathbf{x}$:

$$\mathbf{F}_{\pm} = \mp \frac{\mathbf{v}}{2^{5/2}(\mathbf{w} \cdot \mathbf{x})^{3/2}}, \quad (12)$$

with complex vectors

$$\begin{aligned} \mathbf{v} &= \alpha_1 \mathbf{a}_2 \times \mathbf{a}_3 + \alpha_2 \mathbf{a}_3 \times \mathbf{a}_1 + \alpha_3 \mathbf{a}_1 \times \mathbf{a}_2 \\ \mathbf{w} &= \alpha_1 \mathbf{a}_1 + \alpha_2 \mathbf{a}_2 + \alpha_3 \mathbf{a}_3, \end{aligned} \quad (13)$$

with $\alpha_1^2 + \alpha_2^2 + \alpha_3^2 = 0$. The easily proven result $\mathbf{v} \cdot \mathbf{w} = 0$ guarantees that $\nabla \cdot \mathbf{F}_{\pm} = 0$.

The condition $\mathbf{w} \cdot \mathbf{x} = 0$ is satisfied locally if $\mathbf{x}$ is orthogonal to *both* $\mathbf{w}_R \equiv \text{Re} \mathbf{w}$ and $\mathbf{w}_I \equiv \text{Im} \mathbf{w}$, showing that the line of generalized degeneracies is locally tangent to $\mathbf{u} = \mathbf{w}_R \times \mathbf{w}_I$. Note that unless it is real, $\mathbf{v}$ will

not be parallel to $\mathbf{u}$. This line can pierce the integration surface S at singular points where $\mathbf{F}_\pm$ will have the asymptotic behavior (12). Since the surface integral is two dimensional, the singularity will be *integrable*.

Suppose now that $\mathbf{x}$ circles a singular line in the plane formed by $\mathbf{w}_R$ and $\mathbf{w}_I$. For example, let $\mathbf{w} \cdot \mathbf{x} = \eta e^{i\theta}$ with η fixed and $-\pi < \theta \leq \pi$ determined by $\mathbf{w}_R \cdot \mathbf{x} = \eta \cos(\theta)$ and $\mathbf{w}_I \cdot \mathbf{x} = \eta \sin(\theta)$. Then $\mathbf{F}_\pm \propto e^{-i3\theta/2}$, so that single valuedness can be maintained only if a *surface of branch cuts* extends out of the singular line. It is straightforward to show that this branch sheet joins $\mathbf{F}_+$ and $\mathbf{F}_-$: tracing the closed path causes E_m and E_n to exchange positions by circling each other in the complex plane. $\mathbf{F}_\pm(\mathbf{X})$ forms a single complex function on a two-sheeted Riemann surface (see Fig. 2 below).

In the Hermitean case one may deform the integration surface to small spheres enclosing the point singularities. In the generic case the deformed surface wraps around those portions of the lines and their branch surfaces that are internal to the original surface.

In order to understand the result of the integration (7) we consider a model problem with the *exact* form $\vec{\alpha} = (\alpha_1^0 + \mathbf{a}_1 \cdot \mathbf{x}, \alpha_2^0 + \mathbf{a}_2 \cdot \mathbf{x}, \alpha_3^0 + \mathbf{a}_3 \cdot \mathbf{x})$, so that a true degeneracy exists at $\mathbf{x} = \mathbf{0}$ when $\vec{\alpha}^0 = (\alpha_1^0, \alpha_2^0, \alpha_3^0) = \mathbf{0}$. Near $\mathbf{0}$ the flux is

$$\mathbf{F}_\pm(\mathbf{x}) = \mp \frac{D_0 \mathbf{x}}{2(\mathbf{S}\mathbf{x} \cdot \mathbf{S}\mathbf{x})^{3/2}} \quad (14)$$

where $\mathbf{S}$ is the 3×3 matrix whose rows are the $\mathbf{a}_i$, and $D_0 = \det \mathbf{S} = \mathbf{a}_1 \cdot \mathbf{a}_2 \times \mathbf{a}_3$.

We suppose that the $\mathbf{a}_i$ are all real, so that $\vec{\alpha}^0$ represents a (generally non-Hermitean) perturbation of an *Hermitean* problem. On transformation to real coordinates $\mathbf{y} = \mathbf{S}\mathbf{x}$, the flux takes the classic Coulomb form $\mathbf{F}_\pm(\mathbf{y}) = \mp \sigma(D_0) \mathbf{y} / 2|\mathbf{y}|^3$, where $\sigma(D_0)$ is the sign of D_0 . It follows immediately that the integrated flux through any surface S enclosing the origin is $\gamma(S) = \mp 2\pi \sigma(D_0)$.

On including the constant shift $\vec{\alpha}^0$, the flux becomes

$$\mathbf{F}_\pm(\mathbf{y}) = \mp \sigma(D_0) \frac{\vec{\alpha}^0 + \mathbf{y}}{2[(\vec{\alpha}^0 + \mathbf{y}) \cdot (\vec{\alpha}^0 + \mathbf{y})]^{3/2}}. \quad (15)$$

For very large $|\mathbf{y}|$ this form reduces to (14). The flux through a surface at infinity remains $\gamma(S)$. Since $\nabla_{\mathbf{y}} \cdot \mathbf{F}_\pm(\mathbf{y}) = 0$, this result holds for any surface that encloses all singularities in $\mathbf{F}_\pm$. These singularities occur at the zeros of $(\vec{\alpha}^0 + \mathbf{y})^2$, i.e., for $|\mathbf{y} + \text{Re}\vec{\alpha}^0|^2 = |\text{Im}\vec{\alpha}^0|^2$ and $\text{Im}\vec{\alpha}^0 \cdot (\mathbf{y} + \text{Re}\vec{\alpha}^0) = 0$, which constitute a circular ring of radius $a = |\text{Im}\vec{\alpha}^0|$ formed by the intersection of the sphere of radius a centered at $-\text{Re}\vec{\alpha}^0$ with the plane normal to $\text{Im}\vec{\alpha}^0$ passing through this same center. In $\mathbf{x}$ -space the ring becomes an ellipse. A ring must be spanned by a branch surface in order to maintain single-valuedness of $\mathbf{F}_\pm$.

We have therefore established that under non-Hermitean perturbations the usual quantized point charges expand into closed loops, with net flux quantized by the original Hermitean Chern number. If a loop

grows to pierce the surface of integration, only a fraction of the total flux will be enclosed, and integer quantization is lost. This fraction depends on the geometry of the branch cut, which is itself arbitrary, except for its end points. Only the *total* flux, integrated over *all* sheets of the Riemann surface, remains quantized [11]. The latter will correspond to the net Chern number of the colliding eigenvalues, which are now topologically “entangled.” The entanglement becomes more pronounced as the non-Hermiteity increases, and may involve more than two eigenvalues that collide with each other sequentially as $\mathbf{X}$ is varied, generating further Riemann sheets.

We illustrate this formal discussion by a specific example, a family of operators in two dimensions:

$$\mathbf{L} = -e^{i\theta} D \nabla^2 + i\mathbf{A}(\mathbf{r}) \cdot \nabla, \quad (16)$$

in which $0 \leq \theta < 2\pi$ controls the degree of non-Hermiteity ($\mathbf{L}$ is Hermitean only for $\theta = 0, \pi$), D is a constant coefficient, and $\mathbf{A}$ is an incompressible vector field: $\nabla \cdot \mathbf{A} = 0$. For $\theta = 0$, this operator corresponds to the random flux problem in the Coulomb gauge [6], with magnetic field $B = \nabla \times \mathbf{A}$, and a particular choice $V = -\mathbf{A}^2/4D$ for the scalar potential. For $\theta = \pi/2$ this operator represents a model of flux lines in a three-dimensional superconductor with extended defects, (where $\mathbf{A}$ corresponds to the horizontal components of the magnetic field [3]), or a two-dimensional passive-scalar transport model, (where $\mathbf{A}$ is the velocity field, and D is the diffusion constant). We set $D = 1$ and implement (16) numerically by discretizing it [4] on a small (4×4) lattice. The components of the vector field $\mathbf{A}$ are initially chosen to be independent random variables chosen uniformly on an interval $[-W, W]$, with $W = 40$. Incompressibility is imposed by subtracting from $\mathbf{A}$ the function ∇a , where $a(\mathbf{r})$ satisfies $\nabla^2 a = \nabla \cdot \mathbf{A}$ with periodic boundary conditions.

Study of random operators like (16) is motivated by questions of localization [2–4,6]. Extended states are distinguished from localized states by their sensitivity to boundary conditions. Let $E_n(\phi_x, \phi_y)$, $\psi_n^{\phi_x, \phi_y}(x, y)$ be the eigenvalues and eigenstates of $\mathbf{L}$ on an $L_x \times L_y$ system with boundary conditions $\psi_n^{\phi_x, \phi_y}(x + L_x, y) = e^{i\phi_x} \psi_n^{\phi_x, \phi_y}(x, y)$, $\psi_n^{\phi_x, \phi_y}(x, y + L_y) = e^{i\phi_y} \psi_n^{\phi_x, \phi_y}(x, y)$. The two-dimensional space (ϕ_x, ϕ_y) has the geometry of a torus that is embedded in a three-dimensional parameter space $\mathbf{X}$. The Berry flux through the torus is the integrand of (9): $F_n(\phi_x, \phi_y) = \langle \partial_{\phi_x} \psi_n | \partial_{\phi_y} \psi_n \rangle - \langle \partial_{\phi_y} \psi_n | \partial_{\phi_x} \psi_n \rangle$, and if the integral of the flux over the surface (the Chern number) is non-zero, then the eigenstate is extended.

Fig. 1 displays the variation of eigenvalues with θ for $\phi_x = \phi_y = 0$. In Fig. 2 we plot the imaginary parts of $F_\pm(\phi_x, \phi_y)$ for the labeled eigenvalues in Fig. 1 and for two values of θ , revealing the emergence of a branch cut and showing $F_\pm$ are different sheets of the same analytic function. The log-log plot exhibits the $3/2$ -law (12) as an endpoint of a branch cut is approached.




FIG. 1. The middle six eigenvalues λ as a function of non-Hermiticity parameter θ for $\phi_x = \phi_y = 0$.

The action corresponding to (16) at $\theta = \pi/2$ is the Wick rotation of the action at $\theta = 0$. A natural question is whether this analytic continuation to imaginary time is accompanied by a smooth variation in properties of eigenstates, or whether a phase transition, e.g., metal to insulator, occurs as a function of θ (see ref. [4]). For finite θ , we have shown that the geometry of eigenvalue degeneracies can change dramatically, but the notion of Chern number is still relevant. Whether the entanglement of eigenvalues can lead to a phase transition in the thermodynamic limit is unknown.

In summary, we extend the notion of Chern number to non-Hermitian operators and find that it remains a useful tool for distinguishing extended and localized states. Entanglement of eigenstates can lead to an intricate multi-sheeted analytic structure for the Berry flux, and the Chern number becomes defined only for the group of entangled states as a whole.




FIG. 2. Plot of imaginary part of the Berry flux $\text{Im}F_{\pm}(\phi_x, \phi_y)$ for two values of θ , in a small part of the Brillouin zone, $[0.35\pi, 0.40\pi] \times [1.24\pi, 1.25\pi]$. The Chern numbers for eigenvalues I and II are +1 and -1 respectively until the first branch cut (not shown) occurs at about $\theta = .021\pi$. As shown on the left, another branch cut (the obvious vertical lines in the two lower panels separating white from black) emerges between $\theta = .025\pi$ and $\theta = .029\pi$. The linear grey scale spans the range $[-1.3, 1.3]$. The log-log plot on the right displays the divergence of the magnitude of the flux against the distance r from an endpoint of the branch cut along the paths (h) and (v); the solid line has slope $-3/2$; deviations from it at small r arise from finite numerical resolution.

case this fails in general because E_m is no longer real, and an initially infinitesimal amount of tunneling may grow exponentially instead of simply oscillating. For our purposes the connection with the adiabatic limit is not necessary. It matters only that parallel transport defined by (3) along an arbitrary curve C be implemented dynamically by *some* choice of $\mathbf{L}(t)$.

-
- [1] *Geometric Phases in Physics*, Edited by A. Shapere and F. Wilczek (World Scientific, Singapore, 1989) contains a comprehensive introduction to the subject.
 - [2] Y. Huo and R.N. Bhatt, Phys. Rev. Lett. **47**, 1546 (1992).
 - [3] N. Hatano and D. R. Nelson, Phys. Rev. Lett. **77**, 570 (1996); N. M. Shnerb and D. R. Nelson, Phys. Rev. Lett. **80**, 5172 (1998).
 - [4] J. Miller and J. Wang, Phys. Rev. Lett. **76**, 1461 (1995); J. Miller, *Non-fermi liquid as passive scalar fluid*, cond-mat/9906398.
 - [5] J.C. Garrison and W.M. Wright, Phys. Lett. A **178**, 177 (1988).
 - [6] See [4] and references therein.
 - [7] In the Hermitean case it can be shown [1] that this same dynamics is generated by $\mathbf{L}(\mathbf{X}(\epsilon t))$, after subtracting the time integral of $E_m(\mathbf{X}(t))$, in the adiabatic limit, $\epsilon \rightarrow 0$ but $\alpha = \epsilon t$ finite. In this limit, tunneling to other eigenstates becomes vanishingly small. In the non-Hermitean

- [8] We assume henceforth that $\mathbf{X}$ is a 3-dimensional vector.
- [9] Since $\gamma_m(S_1)$ and $\gamma_m(S_2)$ differ, $\mathbf{A}_m$ must have a line singularity that emerges from the singularity in $\mathbf{F}_m$ and passes through S_1 or S_2 . This is not a true singularity, but a coordinate singularity, as occurs in polar coordinates at the poles of a sphere, and its position varies with gauge choice. These singularities do not appear in $\mathbf{F}_m$ since it is gauge invariant.
- [10] Let $|-\rangle$ is the eigenvector obeying $\mathbf{L}_{mn}|-\rangle = E|-\rangle$. At a generic degeneracy one has $|+\rangle = |-\rangle$, and there exists an independent generalized eigenvector (also called a cyclic vector) $|g\rangle$ that obeys $\mathbf{L}_{mn}|g\rangle = |-\rangle$ and may be chosen so that $\langle g|\pm\rangle = 0$. For details see an advanced text on linear algebra.
- [11] To prove this, observe that the Berry flux through a branch surface is identical for the two eigenvalues involved, but the surface normals are opposite. Branch surfaces make no contribution and the total integral is conserved and equal to the sum of the Chern numbers before the Chern loop pierced the integration surfaces.